arXiv:cond-mat/0504219v1 [cond-mat.stat-mech] 9 Apr 2005

APS/123-QED

Critical Dynamics of Three-Dimensional Spin Systems with Long-Range Interactions

S.V. Belim*
Omsk State University, 55-a, pr. Mira, Omsk, Russia, 644077 \\

(Dated: February 2, 2008)

A field-theoretic description of critical behavior of Ising systems with long-range interactions is obtained by using the PadeBorel summation technique in the two-loop approximation directly in the three-dimensional space. It is shown that long-range interactions affect the relaxation time of the system.

PACS numbers: 64.60.-i

It was shown in [1] that effects due to long-range interaction are essential for the critical behavior of Ising systems. The renormalization-group approach to spin systems with long-range interactions developed in [2] directly in the three-dimensional space made it possible to calculate the static critical exponents in the two-loop approximation. However, analogous calculations of critical dynamics have never been performed for these systems.

In this paper, a field-theoretic description of critical behavior of homogeneous systems with long-range interactions is developed directly for $D = 3$ in the two-loop approximation. The model under analysis is the classical spin system with exchange integral depending on the distance between spins. The corresponding Hamiltonian is

$$H = \frac{1}{2}\sum_{ij} J(|r_i - r_j|)S_i S_j, \qquad (1)$$

where S_i is a spin variable and $J(|r_i r_j|)$ is the exchange integral. This model is thermodynamically equivalent to the $O(n)$-symmetric GinzburgLandauWilson model defined by the effective Hamiltonian

$$H = \int d^D q \Big\{ \frac{1}{2}(\tau_0 + q^\sigma)\varphi^2 + u_0\varphi^4 \Big\}, \qquad (2)$$

where φ is a fluctuating order parameter, D is the space dimension, $\tau_0 \sim |T - T_c|$ (T_c is the critical temperature), and u_0 is a positive constant. Critical behavior strongly depends on the parameter a characterizing the interaction as a function of distance. It was shown in [3] that long-range interaction is essential when $0 < \sigma < 2$, whereas systems with $\sigma = 2$ exhibit critical behavior characteristic of short-range interactions. For this reason, the analysis that follows is restricted to the case of $0 < \sigma < 2$.

Relaxational dynamics of spin systems near the critical temperature can be described by a Langevin-type equation for the order parameter:

$$\frac{\partial\varphi}{\partial t} = -\lambda_0 \frac{\delta H}{\delta\varphi} + \eta + \lambda_0 \mathbf{h}, \qquad (3)$$

where λ_0 is a kinetic coefficient, $\eta(x,t)$ is a gaussian random force (representing the effect of a heat reservoir) defined by the probability distribution

$$P_\eta = A_\eta \exp\left[-(4\lambda_0)^{-1} \int d^d x\, dt\, \eta^2(x,t) \right] \qquad (4)$$

with a normalization factor A_η, and $\mathbf{h}(t)$ is an external field thermodynamically conjugate to the order parameter. The temporal correlation function $G(x,t)$ of the order-parameter field can be found by solving Eq. (3) with $H[\varphi]$ given by (2) for $\varphi[\eta, \mathbf{h}]$ averaging the result over P_η, and retaining the component linear in $\mathbf{h}(0)$

$$G(x,t) = \frac{\delta}{\delta \mathbf{h}(0)}[\langle\varphi(x,t)\rangle]|_{h=0}, \qquad (5)$$

where

$$[\langle\varphi(x,t)\rangle] = B^{-1}\int D\{\eta\}\varphi(x,t)P_\eta, \qquad (6)$$

$$B = \int D\{\eta\}P_\eta. \qquad (7)$$

When applying the standard renormalization-group procedure to this dynamical model, one has to deal with substantial difficulties. However, it was shown in [4] that the model of critical dynamics in homogeneous systems without long-range interaction based on a Langevin-type equation is equivalent to that described by the standard Lagrangian [5]

$$L = \int d^d x\, dt \left\{ \lambda_0^{-1}\varphi^2 + i\varphi^* \left(\lambda_0^{-1}\frac{\partial\varphi}{\partial t} + \frac{\delta H}{\delta\varphi} \right) \right\}, \qquad (8)$$

where φ^* denotes an auxiliary field. The corresponding correlation function $G(x,t)$ of the order parameter is defined for a homogeneous system as

$$\begin{aligned} G(x,t) &= \langle\varphi(0,0)\varphi(x,t)\rangle \qquad (9) \\ &= \Omega^{-1}\int D\{\varphi\}D\{\varphi^*\}\varphi(0,0)\varphi(x,t)\exp(-L[\varphi,\varphi^*]), \end{aligned}$$

where

$$\Omega = \int D\{\varphi\}D\{\varphi^*\}\exp(-L[\varphi,\varphi^*]). \qquad (10)$$

---

*Electronic address: belim@univer.omsk.su

Instead of dealing with the correlation function, it is reasonable to invoke the Feynman diagram technique and represent the corresponding vertex in the two-loop approximation as

$$\Gamma^{(2)}(k,\omega;\tau_0,u_0,\lambda_0) = \tau_0 + k^\sigma - \frac{i\omega}{\lambda_0} - 96{u_0}^2 D_0,$$
$$D_0 = \frac{3}{4}\int \frac{d^Dq d^Dp}{(1+|\vec{q}|^\sigma)(1+|\vec{p}|^\sigma)} \qquad (11)$$
$$\cdot\frac{1}{(3+|\vec{q}|^\sigma+|\vec{p}|^\sigma+|\vec{p}+\vec{q}|^\sigma-i\omega/\lambda)}$$

The next step in the field-theoretic approach is the calculation of the scaling functions β, γ_τ, γ_φ and γ_λ in the renormalization-group differential equation for vertexes

$$\left[\mu\frac{\partial}{\partial\mu} + \beta\frac{\partial}{\partial u} - \gamma_\tau\tau\frac{\partial}{\partial\tau} + \gamma_\lambda\lambda\frac{\partial}{\partial\lambda} - \frac{m}{2}\gamma_\varphi\right]$$
$$\times\Gamma^{(m)}(k,\omega;\tau,u,\lambda,\mu) = 0, \qquad (12)$$

where the scaling parameter μ is introduced to change to dimensionless variables.

Further analysis requires the use of the function β and the dynamic scaling function γ_λ.

An expression for β in the two-loop approximation were obtained in [2]:

$$\beta = -(4-D)\Big[1 - 36uJ_0 + 1728\Big(2J_1 - J_0^2 - \frac{2}{9}G\Big)u^2)\Big],$$
$$J_1 = \int\frac{d^Dq d^Dp}{(1+|\vec{q}|^\sigma)^2(1+|\vec{p}|^\sigma)(1+|q^2+p^2+2\vec{p}\vec{q}|^{\sigma/2})},$$
$$J_0 = \int\frac{d^Dq}{(1+|\vec{q}|^\sigma)^2},$$
$$G = -\frac{\partial}{\partial|\vec{k}|^\sigma}\int\frac{d^Dq d^Dp}{(1+|q^2+k^2+2\vec{k}\vec{q}|^{\sigma/2})(1+|\vec{p}|^\sigma)}$$
$$\cdot\frac{1}{(1+|q^2+p^2+2\vec{p}\vec{q}|^{\sigma/2})}$$

The function γ_λ calculated in the two-loop approximation is

$$\gamma_\lambda = (2\sigma - D)2(D' - G)u^2, \qquad (13)$$
$$D' = \frac{\partial D_0}{\partial(-i\omega/\lambda)}|_{k=0,\omega=0}.$$

Defining the effective interaction vertex $v = u\cdot J_0$ one obtains the following expressions for β and γ_λ:

$$\beta = -(2\sigma-D)\Big[1 - 36v + 1728\Big(2\widetilde{J_1} - 1 - \frac{2}{9}\widetilde{G}\Big)v^2)\Big],$$
$$\gamma_\lambda = (2\sigma-D)96(\widetilde{D}-\widetilde{G})v^2, \qquad (14)$$
$$\widetilde{J_1} = \frac{J_1}{J_0^2} \quad \widetilde{G} = \frac{G}{J_0^2} \quad \widetilde{D} = \frac{D'}{J_0^2}.$$

This redefinition is meaningful for $\sigma \le D/2$. In this case, J_0, J_1, G and D' are divergent functions. Introducing the cutoff parameter Λ, we obtain finite expressions for the ratios

$$\frac{J_1}{J_0^2} = \Big[\int_0^\Lambda\int_0^\Lambda d^Dqd^Dp/((1+|\vec{q}|^\sigma)^2(1+|\vec{p}|^\sigma)$$
$$(1+|q^2+p^2+2\vec{p}\vec{q}|^\sigma))\Big]\Big/\Big[\int_0^\Lambda d^Dq/(1+|\vec{q}|^\sigma)^2\Big]^2,$$
$$\frac{G}{J_0^2} = \Big[-\frac{\partial}{(\partial|\vec{k}|^\sigma)}\int_0^\Lambda\int_0^\Lambda d^Dqd^Dp/((1+|\vec{p}|^\sigma)$$
$$(1+|q^2+k^2+2\vec{k}\vec{q}|^\sigma)(1+|q^2+p^2+2\vec{p}\vec{q}|^\sigma))\Big]$$
$$\Big/\Big[\int_0^\Lambda d^Dq/(1+|\vec{q}|^\sigma)^2\Big]^2, \qquad (15)$$
$$\frac{D'}{J_0^2} = \frac{3}{4}\Big[\int d^Dqd^Dp/((1+|\vec{q}|^\sigma)(1+|\vec{p}|^\sigma)$$
$$(3+|\vec{q}|^\sigma+|\vec{p}|^\sigma+|\vec{p}+\vec{q}|^\sigma)^2)\Big]\Big/\Big[\int_0^\Lambda d^Dq/(1+|\vec{q}|^\sigma)^2\Big]^2.$$

as $\Lambda \to \infty$.

The integrals are performed numerically. For $\sigma \le D/2$, a sequence of J_1/J_0^2 and G/J_0^2 corresponding to various values of Λ is calculated and extrapolated to infinity.

Critical behavior is completely determined by the stable fixed points of the renormalization group (RG) transformation. These points can be found from the condition

$$\beta(v^*) = 0. \qquad (16)$$

The effective interaction vertexes were evaluated at the stable fixed points of the RG transformation in [2].

The dynamic critical exponent z characterizing critical slowing-down of relaxation is determined by substituting the effective charges at a fixed point into the scaling function :

$$z = 2 + \gamma_\lambda. \qquad (17)$$

The table shows the stable fixed points of the RG transformation and the values of the dynamic critical exponent for $1.5 \le \sigma \le 1.9$. When $0 < \sigma < 1.5$, the only fixed point is the unstable gaussian one, $v^* = 0$.

A comparison of the present results with the value $z = 2.017$ of the dynamic critical exponent for three-dimensional systems with short-range interactions obtained in [6] demonstrates the essential role played by long-range interactions in critical dynamics of spin systems. In particular, the system's relaxation time increases according to the scaling $t \sim |T - T_c|^{\nu z}$, where ν is the critical exponent characterizing the increase in the correlation radius near a critical point. In three-dimensional systems with long-range interaction, both critical dynamics and static behavior become increasingly gaussian as the long-range interaction parameter σ decreases. When $a \le 1.8$, critical behavior is virtually gaussian.

The work is supported by Russian Foundation for Basic Research N 04-02-16002.

Table 1.

a	v^*	z
1.5	0.015151	2.000072
1.6	0.015974	2.000180
1.7	0.020485	2.000777
1.8	0.023230	2.001529
1.9	0.042067	2.006628

[1] E. Luijten and H. Mebingfeld, Phys. Rev. Lett. 86, 5305 (2001).

[2] S. V. Belim, Pisma Zh. Eksp. Teor. Fiz. 77, 118 (2003) [JETP Lett. 77, 112 (2003)].

[3] M. E. Fisher, S.-K. Ma, and B. G. Nickel, Phys. Rev. Lett. 29, 917 (1972).

[4] C. De Dominicis, Lett. Nuovo Cimento 12, 567 (1975).

[5] E. Brezin, J. C. LeGuillon, and J. Zinn-Justin, Phys. Rev. D 8, 434 (1973); Phys. Rev. D 8, 2418 (1973).

[6] V. V. Prudnikov, S. V. Belim, A. V. Ivanov, et al., Zh. Eksp. Teor. Fiz. 114, 972 (1998) [JETP 87, 527 (1998)].